

Complimentary and Alternative Treatments for Hepatitis C

National Center for Complimentary and Alternative Medicine

www.nccam.nih

Hepatitis C and Complimentary and Alternative Medicine

Key Points

What is hepatitis C?

What does conventional treatment for chronic hepatitis C consist of?

Why do people use CAM for hepatitis C?

How commonly do people with hepatitis C use CAM therapies, and what do they use?

What CAM therapies are discussed in this Research Report?

What is known from scientific evidence about CAM modalities for hepatitis C?

What should I do to take care of myself if I have hepatitis C?

Scientific Research Findings: Selected CAM Treatments for Hepatitis C

For More Information

Sources

Appendix

Introduction

Hepatitis C is a disease of the liver that is caused by the hepatitis C virus. The disease occurs in acute and chronic forms; symptoms can range from mild (or even no symptoms) to severe. There are conventional medical treatments available for hepatitis C, but some patients also try complementary and alternative medicine (CAM).^a This Research Report answers some frequently asked questions on hepatitis C and CAM, reviews findings from scientific research on some dietary supplements that have been used as CAM treatments for hepatitis C (milk thistle, licorice root, ginseng, thymus extract, schisandra, and colloidal silver), and suggests sources for further information.

^aConventional medicine is medicine as practiced by holders of M.D. (medical doctor) or D.O. (doctor of

osteopathy) degrees and their allied health professionals, such as physical therapists, psychologists, and registered nurses. Other terms for conventional medicine include allopathy; Western, mainstream, orthodox, and regular medicine; and biomedicine. Some conventional medical practitioners are also practitioners of CAM. CAM, as defined by NCCAM, is a group of diverse medical and health care systems, practices, and products that are not presently considered to be part of conventional medicine.

Key Points

Conventional medical treatment (consisting of a combination drug regimen) for hepatitis C has shown sustained benefit in approximately 55 percent of patients.

Some of the reasons hepatitis C patients try CAM are that they find conventional drug treatment difficult to tolerate or they do not experience a sustained response to treatment.

No CAM treatment has yet been proven safe and effective for treating hepatitis C.

There are many CAM treatments for which benefits for health are claimed. However, it is important to find out what scientific studies have been done on the safety and effectiveness of the CAM treatment in which you are interested. Clinical trials^b are needed of CAM therapies that may show some potential for benefit for hepatitis C, such as milk thistle. The National Center for Complementary and Alternative Medicine (NCCAM) is sponsoring a clinical trial of milk thistle.

It is important to inform all of your health care providers about any therapy that you are currently using or considering, including any dietary supplements. This is to help ensure a safe and coordinated course of care.

What is hepatitis C?

Hepatitis C is a communicable (contagious) disease of the liver caused by the hepatitis C virus (HCV).^c The liver, the largest organ in the body, is found behind the ribs on the right side of the abdomen. It has many important functions, including removing harmful material from the blood and converting food into substances needed for life and growth. The term "hepatitis" means inflammation of the liver. There are other viruses in the hepatitis family (such as hepatitis A and hepatitis B), but HCV is not related to them.

Quick Facts About Hepatitis C

Hepatitis C is the most common bloodborne infection in the United States. About 35,000 new cases

are diagnosed in the United States each year.

Hepatitis C is transmitted primarily when an infected person's blood comes into contact with the blood of a noninfected person.

People who are at the highest risk for HCV infection are those who have used or experimented with injection drugs; received a blood transfusion, blood product, or organ transplant before July 1992; worked in health care and had a needlestick accident involving HCV-infected blood; or had multiple sex partners.

A risk exists but is low (1 to 5 percent) for babies born to a mother with hepatitis C and for people who are in a monogamous sexual relationship with someone with hepatitis C; who have had other sexually transmitted diseases; who have had tattooing or body piercing done with unsterilized tools; or who have used cocaine intranasally (i.e., "snorted" it).

Hepatitis C is not spread through sneezing, coughing, kissing, hugging, food or water, or casual contact.

People who are newly infected have what is called acute hepatitis C. For about 15 to 40 percent of this group, the infection is short-term, goes away, and does not return. Others develop chronic (or long-lasting) hepatitis C, in which the virus stays in the liver, replicates itself, and injures the liver over time.

Among people with chronic hepatitis C, most show no symptoms for up to 20 to 30 years; some have mild symptoms; and some have more serious symptoms.

Chronic hepatitis C can cause liver disease, cirrhosis (scarring of the liver), liver cancer, and liver failure. However, persons who have been diagnosed with hepatitis C need to know that serious illness or death from the disease is by no means inevitable--especially if they take proper care of themselves and get the health care they need.

What does conventional treatment for chronic hepatitis C consist of?

People who have a mild case of hepatitis C may only need to manage it by visiting their doctor regularly and following their doctor's recommendations--such as eating a nutritious diet, avoiding alcohol

(because of its impact on the liver), and getting regular exercise.

For people with more severe hepatitis C, however, drug therapy may be needed. A drug called interferon is the mainstay of conventional treatment. Interferon is often combined with an antiviral (virus-fighting) drug called ribavirin. Such combination therapies are usually taken for 6 months to 1 year. Approximately 55 percent of patients treated with the combination of interferon and ribavirin for 1 year will achieve a sustained response (that is, a sustained benefit from treatment).¹ If a patient does not achieve a sustained response, his doctor may decide whether another course of treatment (re-treatment) is appropriate.

Combination regimens benefit many patients. However, their side effects can be difficult for some patients to tolerate. These side effects can include flu-like symptoms (such as body aches, fever, chills, and fatigue); nausea and other gastrointestinal problems; hair loss; emotional changes; skin reactions; and, in more severe cases, depression, organ damage, blood conditions, and other problems.

Why do people use CAM for hepatitis C?

There are various reasons why people use CAM for hepatitis C, including:

They have not had a response to initial treatment or to re-treatment with drugs.

They are not willing to have drug treatment or continue it--for example, because of the side effects or length of treatment.

They would like to support their body's fight against damage by hepatitis C, and they hear of benefits claimed for some CAM treatments--such as "strengthens the immune system" or "cleanses or rejuvenates the liver" (or other organs).

They are experiencing problems from other diseases and conditions that can be caused by or worsened by hepatitis C.

They are not satisfied with their conventional medical treatment.

How commonly do people with hepatitis C use CAM therapies, and what do they use?

While there have been no surveys yet on the use of CAM by persons with hepatitis C specifically, there is some data from a survey published in 2002 on the use of CAM by persons who have chronic liver diseases (such as hepatitis, liver cancer, alcoholic liver disease, or cirrhosis).² This survey of 989 patients

being treated for various liver diseases at six clinics in the United States found that 39 percent used some form of "alternative therapy." The therapy they used the most was herbals or botanicals^d (21 percent). However, the herbals and botanicals were used for reasons besides liver disease, such as depression. Thirteen percent of all survey participants used herbals or botanicals specifically for their liver disease, and they used only milk thistle (12 percent) or licorice root (1 percent). The other most commonly used CAM therapies were self-prayer^e (18 percent), and (from 6 to 9 percent each) relaxation, megavitamins, massage, chiropractic, and spiritual healing.²

^dHerbs are plants or plant parts valued for their flavor, scent, and/or therapeutic properties. "Herbals" and "botanicals" are synonyms and mean herbal and botanical products.

^eSelf-prayer is when an individual prays for himself. It can be contrasted with intercessory prayer, in which an individual prays for others.

What CAM therapies are discussed in this Research Report?

There is a range of medical concerns associated with hepatitis C, and the number of CAM therapies that are tried is large.^f Therefore, it is beyond the scope of this Research Report to discuss all possible CAM therapies used for hepatitis C. The report focuses on a number of dietary supplements that are used: milk thistle, licorice root, ginseng, thymus extract, schisandra, and colloidal silver.

About Dietary Supplements

Dietary supplements were defined in a law passed by Congress in 1994. A dietary supplement must meet all of the following conditions:

It is a product (other than tobacco) intended to supplement the diet, which contains one or more of the following: vitamins, minerals, herbs or other botanicals, amino acids, or any combination of the above ingredients.

It is intended to be taken in tablet, capsule, powder, softgel, gelcap, or liquid form.

It is not represented for use as a conventional food or as a sole item of a meal or the diet.

It is labeled as being a dietary supplement.

What is known from the scientific evidence about CAM

modalities for hepatitis C?

No CAM treatment has been scientifically proven to successfully treat hepatitis C.

Authors who have done recent analyses of the scientific work have found some results that are intriguing and even promising, but they have noted that more research--especially in the form of controlled clinical trials--is needed before firm conclusions can be drawn.

The authors of a 2003 systematic review of medicinal herbs for hepatitis C concluded that there is not enough evidence to support using herbs to treat the disease. This team identified 13 clinical trials that were of sufficient quality for them to analyze. Compared to placebo,^h they found that none of the herbs tested showed effects on liver enzymes or reduced the amount of HCV in the bloodstream, except for milk thistle, which did show a significant reduction of liver enzymes in one trial.³

Two general reviews from 2000 that covered a variety of CAM modalities for hepatitis C concluded that conventional therapies are the only scientifically proven treatments for the disease.^{4,5}

NIH released a Consensus Statement in 2002 on the management of hepatitis C.ⁱ This assessment by a panel of medical and scientific experts found that "alternative and nontraditional medicines" should be studied.

^hA placebo is designed to resemble as much as possible the treatment being studied in a clinical trial, except that the placebo is inactive. An example of a placebo is a pill containing sugar instead of the drug or other substance being studied. By giving one group of participants a placebo and the other group the active treatment, the researchers can compare how the two groups respond and get a truer picture of the active treatment's effects. In recent years, the definition of placebo has been expanded to include other things that could have an effect on the results of health care, such as how a patient and a health care provider interact, how a patient feels about receiving the care, and what he or she expects to happen from the care.

What should I do to take care of myself if I have hepatitis C?

Make sure you have received an accurate diagnosis. Hepatitis C can be diagnosed reliably only through sophisticated blood tests used in conventional medicine.

See your health care provider regularly.

Discuss treatment options with your provider. Ask any questions you have to make sure you understand any treatment and possible side effects. Follow her recommendations for any changes to your diet and/or lifestyle.

Tell your provider about any herbal supplements, other dietary supplements, or medications (whether prescription or over-the-counter) that you are using or considering. This is important for your safety. Even if your provider does not know about the actions or interactions of an herbal supplement or other CAM treatment, he can access the most current medical guidance.

Get vaccinated against hepatitis A and B. Infection with hepatitis C does not prevent a person from becoming infected with other types of hepatitis; if this happens, it can be serious, even life-threatening.

Be an informed consumer. Seek high-quality, science-based information on any CAM modality that you are using or considering. There is free information from NCCAM, the National Library of Medicine, and other Federal sources to help you distinguish science-based information from other types, including word-of-mouth and manufacturers' claims.

If you decide to try herbal supplements, do so with care. (See the NCCAM fact sheet "Herbal Supplements: Consider Safety, Too.")

If you would like to find out about clinical trials of treatments for hepatitis C, go to www.clinicaltrials.gov or contact the NCCAM Clearinghouse.

Scientific Research Findings: Selected CAM Treatments for Hepatitis C

This section describes six CAM therapies that people have used to treat hepatitis C.

There are different types of review articles: In a general review, a broad picture of the scientific studies and evidence available on a particular topic is presented. In a systematic review, data from a set of studies on a particular question or topic are collected, analyzed, and critically reviewed. A meta-analysis uses statistical techniques to analyze results from a collection of individual studies.

Milk Thistle

Milk thistle (scientific name *Silybum marianum*) is a plant from the aster family. The active extract of milk thistle believed to be responsible for the herb's medicinal qualities is silymarin, found in the fruit.⁶ Milk thistle has been used in Europe as a treatment for liver disease and jaundice since the 16th century.⁷

Summary of the research findings

The results of scientific studies to date do not definitively find that milk thistle is beneficial in treating hepatitis C in humans.

Studies in laboratory animals suggest that silymarin may have various benefits to the liver, such as promoting the growth of certain types of liver cells, having a protective effect upon liver cells, fighting a chemical process called oxidation that can damage cells, and inhibiting inflammation.⁷⁻¹⁴ However, in

some cases, a consistent pattern of benefit was not seen, and these studies did not specifically examine the effects of silymarin on hepatitis C.

There have been some studies on silymarin or milk thistle in humans. These studies have generally been small and on liver diseases rather than on hepatitis C infection specifically, and the results have been contradictory (with some positive and some negative).¹⁵⁻¹⁷ A review and a meta-analysis published in 2001 on silymarin in the treatment of liver diseases found it to be generally safe, but contained no firm conclusions with regard to its use to treat viral hepatitis.^{18,19} A 2002 systematic review on milk thistle for liver disease found "no reduction in mortality (frequency of death as an outcome), in improvements in histology (tissue studies) observed through liver biopsy, or in biochemical markers of liver function" and that the data was too limited to support recommending milk thistle for treatment of liver disease.²⁰

To obtain more extensive and reliable data, NCCAM is sponsoring a clinical trial on the use of milk thistle for hepatitis C.

Side effects and other risks

Milk thistle is generally well-tolerated and has shown few side effects in clinical trials. It can cause a laxative effect; less common effects include nausea, diarrhea, abdominal bloating, fullness, and pain. Milk thistle can produce allergic reactions, which tend to be more common among people who are allergic to plants in the same family (e.g., ragweed, chrysanthemum, marigold, and daisy).

Licorice Root

Licorice root is the peeled or unpeeled dried root of the licorice plant (*Glycyrrhiza glabra*). The primary active component of licorice root is a substance called glycyrrhizin. Licorice root has been in use in China since the second and third century B.C. and in the West since Egyptian, Greek, and Roman times.²¹

Summary of the research findings

Laboratory studies of glycyrrhizin in cell cultures suggest that it may have antiviral properties.²¹

In a review of several randomized controlled trials, researchers reported that glycyrrhizin has potential for reducing long-term complications in chronic hepatitis C in those patients who may not respond to interferon.²² Several of the trials reviewed indicated improvements in liver tissue damaged by hepatitis. Some also showed improvements in how well the liver did its job after treatment.

A 1997 study and a 2002 review suggest that long-term administration of glycyrrhizin might prevent liver cancer in patients with chronic hepatitis C.^{23,24}

The use of glycyrrhizin as a complementary therapy (i.e., used in addition to conventional interferon therapy) has been studied, but no significant benefit has been found.^{25,26}

Recent clinical trials have shown that taking glycyrrhizin lowers the levels of liver enzymes (increased

levels of certain liver enzymes indicate liver damage or inflammation). However, taking the herb did not reduce the amount of HCV in patients' blood, a critical indicator of the long-term progress of the infection.²⁷⁻²⁹

Side effects and possible risks

Taking licorice over a prolonged period of time can lead to potentially serious side effects, including high blood pressure, salt and water retention, swelling, depletion of potassium, headache, and/or sluggishness.³⁰ Glycyrrhizin can worsen ascites, the accumulation of fluid in the abdominal cavity, a condition that can be caused by cirrhosis.³¹ The herb also can interact with certain drugs, such as diuretics, digitalis, antiarrhythmic agents, and corticosteroids.

Ginseng

The herb ginseng comes in two types: American ginseng (*Panax quinquefolius*) and Asian ginseng (*Panax ginseng*). Among the Asian forms of ginseng are Chinese, Japanese, and Korean ginseng. (So-called "Siberian ginseng" is not a true ginseng.) Ginseng has been used for thousands of years in Asia. It is usually used with the belief that it will boost the immune system and increase stamina; such properties are thought to be more useful for the elderly and those recovering from illness.³²

Summary of the research findings

The research on ginseng that has been done to date has been primarily in animal models and human tissue in the laboratory. Some beneficial effects of ginseng on the liver were seen in these studies. Researchers concluded that ginseng may also help strengthen glandular systems and the ability to resist disease.³³⁻³⁶

One study found that ginseng may be helpful for elderly people with liver conditions similar to hepatitis.³⁷

No conclusions can be drawn about the possible usefulness and safety of ginseng as a treatment in people who have hepatitis C, because it has not yet been studied formally in people.

Side effects and possible risks

General adverse (negative) effects of ginseng can include insomnia, headache, nosebleed, nervousness, and vomiting. Prolonged use of caffeine and a high dose of ginseng may be associated with hypertension, which is of particular concern for people with cardiovascular disease or diabetes. In addition, people with diabetes who use insulin should be aware that ginseng has demonstrated hypoglycemic effects (lowering of the blood sugar). Ginseng has been shown in laboratory studies to inhibit grouping of platelets in the blood, increasing bleeding risk. Because of this, using ginseng along with NSAIDs (non-steroidal anti-inflammatory drugs), such as aspirin or ibuprofen, should be discussed with your health care provider.³²

Thymus Extract

The thymus is a gland that is involved in the regulation of the body's immune response. Thymus extract

products consist of peptides taken from the thymus glands of cows or calves and are sold as dietary supplements. Often, these products carry claims of boosting immune system functioning to combat diseases, such as hepatitis C. These over-the-counter supplements should not be confused with the prescription drug thymosin alpha-1.

Summary of the research findings

There has been little testing of bovine thymus extract for treatment of hepatitis C. A small clinical trial of a product called Complete Thymic Formula, which contains bovine thymus extracts along with vitamins, herbs, minerals, and enzymes, did not find the product beneficial for hepatitis C patients who had not responded previously to interferon therapy.³⁸ However, this small study does not provide sufficient evidence to draw firm conclusions about either Complete Thymic Formula or thymus extracts in general.

Side effects and possible risks

In the study of Complete Thymic Formula, one adverse event was reported: a patient developed thrombocytopenia, a drop in the number of platelet cells in the blood; the patient recovered after treatment was stopped.³⁸ In general, no adverse effects from thymus extracts have been reported. However, since thymus extracts are derived from animals, there can be concern related to possible contamination from diseased animal parts.^k Accordingly, people on immunosuppressive drugs or who have suppressed immune systems, such as transplant recipients or persons with HIV/AIDS, should use caution about thymus extracts and consult with their health care provider.

Schisandra

Schisandra is a plant that has been used (through extracts from its fruit) in traditional Chinese medicine and in Kampo, traditional Japanese medicine. There are several species, including *Schisandra chinensis*, native to northeastern China and Korea, and *Schisandra sphenanthera*, native to China.

Summary of the research findings

Research has primarily focused on the various lignans (a class of plant nutrients) and essential oils in the dried fruit of schisandra.³⁹ Major constituents include the lignans gomisin A, schizandrins and schizandrol, vitamins C and E, and others.

Studies of the effects of schisandra in the liver have mostly been in animal models. These studies have suggested that extracts of the fruit have a liver-protective effect, a helpful effect on some liver enzymes, and an antioxidant effect.^{l,39,40}

Schisandra is also used in herbal formulas. For example, an herbal medicine called TJ-108 (Ninjin-yomei-to is one of its Japanese names) used in Kampo has schisandra fruit among its herbal components. In one very small study, TJ-108 was compared with two other Kampo herbal formulas for effects in 37 patients who had chronic hepatitis C and had been treated before with interferon.⁴¹ The findings were that TJ-108 may have antiviral properties, which the authors attributed to schisandra fruit and its lignan gomisin A.^{7,41} These findings need to be interpreted with caution because of the study's small size and

because use of an herbal formula, not schisandra alone, was evaluated; herbal formulas contain many ingredients that could cause a variety of effects.

There are no reports on the safety and effectiveness of using schisandra alone for treatment of hepatitis C in humans in the sources reviewed for this report.

Side effects and other risks

Schisandra is considered generally safe. In some people, however, it may cause heartburn, acid indigestion, decreased appetite, stomach pain, or allergic skin rashes.

^lAntioxidants are substances (such as vitamin E) that help prevent oxygen from reacting with other chemicals in cells (oxidation), a process that can have negative effects.

Colloidal Silver

Silver is a metallic element that is mined as a precious metal. People are exposed to silver, usually in tiny amounts, through their environment, drinking water, food, and possibly work or hobbies. Colloidal silver supplements consist of tiny silver particles suspended in a liquid base. They are often marketed with a variety of unproven health claims, including for immunity, diabetes, cancer, and AIDS.

Summary of the research findings

Silver has had some medicinal uses going back for centuries. However, more modern and less toxic drugs have eliminated the vast majority of these uses. Reviews in the scientific literature on colloidal silver have concluded that^{42,43}:

Silver has no known function in the body.

Silver is not an essential mineral supplement or a cure-all and should not be promoted as such.

Claims that there can be a "deficiency" of silver in the body and that such a deficiency can lead to disease are unfounded.

Claims made about the effectiveness of colloidal silver products for numerous diseases are unsupported scientifically.

Colloidal silver products can have serious side effects (discussed below).

Laboratory analysis has shown that the amounts of silver in these supplements vary greatly, which can pose risks to the consumer.

Side effects and other risks

Animal studies have shown that silver builds up in the tissues of the body. In humans, this accumulation can have a serious side effect called argyria, a bluish-gray discoloration of the body, especially of the skin, other organs, deep tissues, nails, and gums. How this happens is not fully known, but silver-protein complexes are thought to deposit in the skin and then be processed by sunlight (similar to traditional

photography).^{44,45} Argyria is not treatable or reversible. Other possible problems include neurologic problems (such as seizures), kidney damage, stomach distress, headaches, fatigue, and skin irritation. Colloidal silver may interfere with the body's absorption of the following drugs: penicillamine, quinolones, tetracyclines, and thyroxine. For more information about colloidal silver, see the NCCAM fact sheet "[Colloidal Silver Products](#)."

References

- National Institute of Diabetes and Digestive and Kidney Diseases. *Chronic Hepatitis C: Disease Management*. National Institute of Diabetes and Digestive and Kidney Diseases Web site. Accessed at digestive.niddk.nih.gov/ddiseases/pubs/viralhepatitis/index.htm on September 3, 2003.
- Strader DB, Bacon BR, Lindsay KL, et al. Use of complementary and alternative medicine in patients with liver disease. *The American Journal of Gastroenterology*. 2002;97(9):2391-2397.
- Liu J, Manheimer E, Tsutani K, et al. Medicinal herbs for hepatitis C virus infection: a Cochrane hepatobiliary systematic review of randomized trials. *The American Journal of Gastroenterology*. 2003;98(3):538-544.
- Kasahara A. Treatment strategies for chronic hepatitis C virus infection. *Journal of Gastroenterology*. 2000;35(6):411-423.
- Sarin SK. Management of hepatitis C: what should we advise about adjunctive therapies, including herbal medicines, for hepatitis C? *Journal of Gastroenterology and Hepatology*. 2000;15(suppl):E164-E171.
- Bean P. The use of alternative medicine in the treatment of hepatitis C. *American Clinical Laboratory*. 2002;21(4):19-21.
- Seeff LB, Lindsay KL, Bacon BR, et al. Complementary and alternative medicine in chronic liver disease. *Hepatology*. 2001;34(3):595-603.
- Flora K, Hahn M, Rosen H, et al. Milk thistle (*Silybum marianum*) for the therapy of liver disease. *The American Journal of Gastroenterology*. 1998;93(2):139-143.
- O'Hara M, Kiefer D, Farrell K, et al. A review of 12 commonly used medicinal herbs. *Archives of Family Medicine*. 1998;7(6):523-536.
- Muriel P, Garciapina T, Perez-Alvarez V, et al. Silymarin protects against paracetamol-induced lipid peroxidation and liver damage. *Journal of Applied Toxicology*. 1992;12(6):439-442.
- Letteron P, Labbe G, Degott C, et al. Mechanism for the protective effects of silymarin against carbon tetrachloride-induced lipid peroxidation and hepatotoxicity in mice: evidence that silymarin acts both as an inhibitor of metabolic activation and as a chain-breaking antioxidant. *Biochemical Pharmacology*.

1990;39(12):2027-2034.

Davila JC, Lenherr A, Acosta D. Protective effect of flavonoids on drug-induced hepatotoxicity in vitro. *Toxicology*. 1989;57(3):267-286.

Fuchs EC, Weyhenmeyer R, Weiner OH. Effects of silibinin and of a synthetic analogue on isolated rat hepatic stellate cells and myofibroblasts. *Arzneimittel-Forschung*. 1997;47(12):1383-1387.

Boigk G, Stroedter L, Herbst H, et al. Silymarin retards collagen accumulation in early and advanced biliary fibrosis secondary to complete bile duct obliteration in rats. *Hepatology*. 1997;26(3):643-649.

Ferenci P, Dragosics B, Dittrich H, et al. Randomized controlled trial of silymarin treatment in patients with cirrhosis of the liver. *Journal of Hepatology*. 1989;9(1):105-113.

Pares A, Planas R, Torres M, et al. Effects of silymarin in alcoholic patients with cirrhosis of the liver: results of a controlled, double-blind, randomized and multicenter trial. I. 1998;28(4):615-621.

Buzzelli G, Moscarella S, Giusti A, et al. A pilot study on the liver protective effect of silybin-phosphatidylcholine complex (IdB1016) in chronic active hepatitis. *International Journal of Clinical Pharmacology, Therapy and Toxicology*. 1993;31(9):456-460.

Wellington K, Jarvis B. Silymarin: a review of its clinical properties in the management of hepatic disorders. *BioDrugs: Clinical Immunotherapeutics, Biopharmaceuticals and Gene Therapy*. 2001;15(7):465-489.

Saller R, Meier R, Brignoli R. The use of silymarin in the treatment of liver diseases. *Drugs*. 2001;61(14):2035-2063.

Jacobs BP, Dennehy C, Ramirez G, et al. Milk thistle for the treatment of liver disease: a systematic review and meta-analysis. *The American Journal of Medicine*. 2002;113(6):506-515.

Shibata S. A drug over the millennia: pharmacognosy, chemistry, and pharmacology of licorice. *Yakugaku Zasshi (Journal of the Pharmaceutical Society of Japan)*. 2000;120(10):849-862.

van Rossum TG, Vulto AG, de Man RA, et al. Review article: glycyrrhizin as a potential treatment for chronic hepatitis C. *Alimentary Pharmacology & Therapeutics*. 1998;12(3):199-205.

Arase Y, Ikeda K, Murashima N, et al. The long term efficacy of glycyrrhizin in chronic hepatitis C patients. *Cancer*. 1997;79(8):1494-1500.

Kumada H. Long-term treatment of chronic hepatitis C with glycyrrhizin [stronger neo-minophagen C (SNMC)] for preventing liver cirrhosis and hepatocellular carcinoma. *Oncology*. 2002;62(suppl 1):94-100.

Abe Y, Ueda T, Kato T, et al. Effectiveness of interferon, glycyrrhizin combination therapy in patients with chronic hepatitis C. *Nippon Rinsho (Japanese Journal of Clinical Medicine)*. 1994;52(7):1817-1822.

Okuno T, Arai K, Shindo M. Efficacy of interferon combined glycyrrhizin therapy in patients with chronic

hepatitis C resistant to interferon therapy. *Nippon Rinsho (Japanese Journal of Clinical Medicine)*. 1994;52(7):1823-1827.

van Rossum TG, Vulto AG, Hop WC, et al. Intravenous glycyrrhizin for the treatment of chronic hepatitis C: a double-blind, randomized, placebo-controlled phase I/II trial. *Journal of Gastroenterology and Hepatology*. 1999;14(11):1093-1099.

Tsubota A, Kumada H, Arase Y, et al. Combined ursodeoxycholic acid and glycyrrhizin therapy for chronic hepatitis C virus infection: a randomized controlled trial in 170 patients. *European Journal of Gastroenterology & Hepatology*. 1999;11(10):1077-1083.

van Rossum TG, Vulto AG, Hop WC, et al. Glycyrrhizin-induced reduction of ALT in European patients with chronic hepatitis C. *The American Journal of Gastroenterology*. 2001;96(8):2432-2437.

Radix glycyrrhizae. In: *WHO Monographs on Selected Medicinal Plants*. Vol. 1. Geneva, Switzerland: World Health Organization; 1999:183-194.

Lewis JH. Licorice for hepatitis C: yum-yum or just ho-hum? *The American Journal of Gastroenterology*. 2001;96(8):2291-2292.

Abebe W. Herbal medication: potential for adverse interactions with analgesic drugs. *Journal of Clinical Pharmacy and Therapeutics*. 2002;27(6):391-401.

Jeong TC, Kim HJ, Park JI, et al. Protective effects of red ginseng saponins against carbon tetrachloride-induced hepatotoxicity in Sprague Dawley rats. *Planta Medica*. 1997;63(2):136-140.

Matsuda H, Samukawa K, Kubo M. Anti-hepatitic activity of ginsenoside Ro. *Planta Medica*. 1991;57(6):523-526.

Nguyen TD, Villard PH, Barlatier A, et al. Panax vietnamensis protects mice against carbon tetrachloride-induced hepatotoxicity without any modification of CYP2E1 gene expression. *Planta Medica*. 2000;66(8):714-719.

Tran QL, Adnyana IK, Tezuka Y, et al. Hepatoprotective effect of majonoside R2, the major saponin from Vietnamese ginseng (*Panax vietnamensis*). *Planta Medica*. 2002;68(5):402-406.

Zuin M, Battezzati PM, Camisasca M, et al. Effects of a preparation containing a standardized ginseng extract combined with trace elements and multivitamins against hepatotoxin-induced chronic liver disease in the elderly. *The Journal of International Medical Research*. 1987;15(5):276-281.

Raymond RS, Fallon MB, Abrams GA. Oral thymic extract for chronic hepatitis C in patients previously treated with interferon: a randomized, double-blind, placebo-controlled trial. *Annals of Internal Medicine*. 1998;129(10):797-800.

Sinclair S. Chinese herbs: a clinical review of Astragalus, Ligusticum, and Schizandrae. *Alternative Medicine Review: A Journal of Clinical Therapeutics*. 1998;3(5):338-344.

Liu GT. Pharmacological actions and clinical use of fructus schizandrae. *Chinese Medical Journal*. 1989;102(10):740-749.

Cyong JC, Kim SM, Iijima K, et al. Clinical and pharmacological studies on liver diseases treated with Kampo herbal medicine. *The American Journal of Chinese Medicine*. 2000;28(3-4):351-360.

Fung MC, Bowen DL. Silver products for medical indications: risk-benefit assessment. *Journal of Toxicology. Clinical Toxicology*. 1996;34(1):119-126.

Gulbranson SH, Hud JA, Hansen RC. Argyria following the use of dietary supplements containing colloidal silver protein. *Cutis*. 2000;66(5):373-374.

White JM, Powell AM, Brady K, et al. Severe generalized argyria secondary to ingestion of colloidal silver protein. *Clinical and Experimental Dermatology*. 2003;28(3):254-256.

Hori K, Martin TG, Rainey P, et al. Believe it or not--silver still poisons! *Veterinary and Human Toxicology*. 2002;44(5):291-292.